

## Bony structure dimensions-fish length relationships of pike (*Esox lucius* L., 1758) in Lake Ladik (Samsun, Turkey)

Okan YAZICIOĞLU<sup>1\*</sup>, Savaş YILMAZ<sup>2</sup>, Mesut ERBAŞARAN<sup>2</sup>,  
Selma UĞURLU<sup>3</sup> and Nazmi POLAT<sup>2</sup>

1. Organic Farming Program, Botanic and Animal Production Department, Technical Vocational Schools of Higher Education, Ahi Evran University, Kırşehir, Turkey.

2. Department of Biology, Faculty of Arts and Science, Ondokuz Mayıs University, Kurupelit, Samsun, Turkey.

3. Department of Biology, Faculty of Arts and Science, Sakarya University, Esentepe Campus, Sakarya, Turkey.

\*Corresponding author, O. Yazicioğlu, Tel: +90 3862804708, E-mail: oknyzcoglu@gmail.com

Received: 11. January 2016 / Accepted: 14. April 2016 / Available online: 26. June 2016 / Printed: June 2017

**Abstract.** The relationships between otolith length (OL), otolith width (OW), otolith weight (OWE), cleithrum length (CL) and fish length (TL) of pike captured from Lake Ladik were evaluated using 11 different mathematical models. Total lengths of the samples ranged from 273 mm to 744 mm. The variations in the right and the left side measurements of the two bony structures were not statistically significant ( $p>0.05$ ). There was significant difference in cleithrum lengths of females and males ( $p<0.05$ ). The relationships between TL versus OL, TL versus OW and TL versus CL were determined as linear, whereas the relationship between TL versus OWE was determined as logarithmic.

**Key words:** *Esox lucius*, otolith biometry, cleithrum length, Lake Ladik, Turkey.


Otoliths or earstones that used for balance and hearing are calcareous structures and located in the inner ears of all teleosts (Campana 1999). There are three pairs of otoliths: sagittae, asteriscus and lapillus (Das 1994). Cleithra, the major bones of the pectoral girdle, are paired structures located on each side of the fish (Scharf et al. 1998). Otoliths and cleithra show differences in size and shape between fish species (Traynor et al. 2010, Yilmaz et al. 2015) and are used to investigate identify, age, and size of prey fishes. This information is very useful for fishery management, prey-predator studies and archaeological investigation (Harvey et al. 2000, Tuset et al. 2008)

Relationships between bony structure dimensions and fish length are commonly used in fisheries science. These mathematical associations enable the back-calculation of fish length in previous ages (Casselman 1990). Such relationships generally show dramatic differences from species to species. Hard parts such as otolith and cleithrum can stay for a long time without being digested in the stomach of piscivorous animals. Thus, the original size of ingested fish is commonly estimated with regression equations that relate body length to the size of a specific hard bone (Aydin et al. 2004, Dietrich et al. 2006). Also, undigested structures found in predator stomach are an important tool for identification of prey species (Tarkan et al. 2007, Mehanna et al. 2016). In addition, bony structures of fish also help paleontologists in

their researches about fish fossils (Jawad et al. 2012).

Otoliths and cleithra have been used in age determination (Laine et al. 1991, Neuman et al. 1994, Faust et al. 2013), species-specific identification (Traynor et al. 2010), and estimation of growth parameters of pike (Rydell et al. 2008). However, the studies on morphometry of these bony structures are very limited for pike (Engstedt et al. 2010). The aim of this study was to determine the best mathematical models defining relationships between otolith and cleithrum dimensions and fish length of pike (*Esox lucius* L., 1758) caught from Lake Ladik (Samsun).

Lake Ladik is between 35° 40'-36° 05' east longitudes and 40° 50'-41° 00' north latitudes. The lake is 5 km long, 2 km wide and 2.5 - 6 m deep with an altitude of 867 m (Yilmaz et al. 2012). It has been classified as a eutrophic and shallow lake (Apaydin Yagci et al. 2015). *Esox lucius* samples were obtained from different parts of the lake between November 2009-October 2010 and February-March 2011. Total lengths of the samples were measured with a precision of  $\pm 1$  mm, samples were weighed with a precision of  $\pm 0.01$  g and sexes were determined through macroscopic examination of gonads. The sagittal otolith and cleithra pairs from each fish were removed, cleaned and preserved dry in labeled envelopes. Sagittal otoliths (OWE) were weighed with a precision of  $\pm 0.0001$  g. Broken and damaged otoliths were not examined. Otolith length (OL) was defined as the farthest distance between the anterior and posterior edges of otolith, and otolith width (OW) was defined as the farthest distance between the dorsal



**Figure 1.** Right otolith (x10) (a) and right cleithrum (x3.2) (b) of pike and measurement lines.

and ventral edges (Fig. 1). OL and OW were measured to the nearest  $\pm 0.001$  mm using Leica Application Suit imaging analysis program on Leica S8APO model stereo microscope. Cleithrum length (CL), which described as the distance from the center of cleithrum to the anterior edge (Casselman 1979) was measured by an electronic caliper of  $\pm 0.01$  mm precision (Fig. 1).

The paired t-test was used for the comparison of the right and left measurements of the two bony structures. Also, this test was used to find out whether the same measurements were different between the sexes or not. The analysis of variance (ANOVA) was applied to determine the best model out of 11 different mathematical models (Linear, Logarithmic, Invers, Quadratic, Cubic, Compound, Power, S, Growth, Exponential and Logistics) defining each relationship (TL-OL, TL-OW, TL-OWE, and TL-CL) with SPSS-16 package program. The model with the lowest standard deviation and the highest regression coefficient ( $r^2$ ) was used to explain the mathematical relationships (Can 2000).


Of the total 210 *Esox lucius* samples caught for the study, 114 were females and 96 were males. While the cleithrum measurements were made for all the samples, otolith measurements were taken from

**Table 1.** The measured values of fish total length, otolith length, and cleithrum length in female, male and all individuals of pike from Lake Ladik (Min: minimum, Max: maximum, SE: standard error, SD: standard deviation).

	Variable	n	Mean	Min	Max	SE	SD
Cleithrum	Female						
	Total length (mm)	114	459.8	273.0	744.0	8.28	88.43
	Cleithrum length (mm)	114	45.64	27.78	73.64	0.83	8.84
	Male						
	Total length (mm)	96	415.2	283.0	584.0	8.55	83.80
	Cleithrum length (mm)	96	41.39	28.41	58.83	0.86	8.44
Otolith	All samples						
	Total length (mm)	210	439.4	273.0	744.0	6.14	88.97
	Cleithrum length (mm)	210	43.70	27.78	73.64	0.61	8.89
	All samples						
	Total length (mm)	134	443.4	273.0	744.0	7.55	87.45
	Otolith length (mm)	134	6.428	4.506	9.958	0.092	1.060
	Otolith width (mm)	134	3.251	2.368	5.030	0.043	0.492
	Otolith weight (g)	134	0.0161	0.0067	0.0516	0.0006	0.0079

**Table 2.** Statistical comparison between right and left side of otolith and cleithrum measurements in pike from Lake Ladik (mm: millimeter, g: grams).

Variable	Area	n	Mean	Min	Max	SE	SD	P
Otolith length (mm)	Left	134	6.417	4.425	9.907	0.094	1.083	$p > 0.05$
	Right	134	6.428	4.506	9.958	0.092	1.060	
Otolith width (mm)	Left	134	3.231	2.270	4.898	0.042	0.485	$p > 0.05$
	Right	134	3.251	2.368	5.030	0.043	0.492	
Otolith weight (g)	Left	134	0.0159	0.0062	0.0508	0.0006	0.0076	$p > 0.05$
	Right	134	0.0161	0.0067	0.0516	0.0006	0.0079	
Cleithrum length (mm)	Left	210	43.44	27.78	73.16	0.62	8.92	$p > 0.05$
	Right	210	43.70	27.78	73.64	0.61	8.89	



**Figure 2.** Relationships between (A) total length-otolith length, (B) total length-otolith width and (C) total length-otolith weight (D) total length-cleithrum length in female and (E) total length-cleithrum length in male of pike from Lake Ladik

only 134 samples.

The total lengths of the fish were between 273-744 mm. Cleithrum lengths were between 27.78-73.64 mm (Table 1). The lengths, widths, and weights of otoliths varied between 4.506-9.958 mm, 2.368-5.030 mm, 0.0067-0.0516 g, respectively (Table 1).

There was no significant difference in the measurements of otolith's length, width, weight, and cleithrum length between the right and the left areas (paired t-test,  $p > 0.05$ ) (Table 2). No significant difference was also observed in the same measurements of females and males (t-test,  $p > 0.05$ ). Thus, the right side measurements were used for regression models regardless of sex difference. On the contrary, since cleithrum length differed between sexes (paired t-test,  $p < 0.05$ ), regressions were generated separately according to female and male.

As a result of the analysis of 11 different mathematical models applied on the otolith data,

it was found that the linear model best explained the relationships between TL-OL and TL-OW, while the logarithmic model was found to be more suitable for the relationship between TL-OWE (Table 2). The  $r^2$  values of the regression equation generated from each measurement of otoliths were same (0.89). This result showed that all three parameters can be used for the back-calculation of fish total length (Fig. 2). A strong linear relationship ( $r^2 = 0.98$ ) was seen between TL-CL of the female and male pikes; approximately 10 times of the cleithrum length was found to be equal to fish total length (Fig. 2).

Since growth of fish can differ between species and the populations of the same species, the relationships between bony structures and fish length can also differ. It is possible to determine the aforementioned relationships in the most correct way through comparing various mathematical models. Gamboa (1991) explained the relationships between fish length-otolith length and fish

length-otolith weight of 11 different fish species in the Pacific shore of California through linear and power models, respectively. Granoderio & Silva (2000) reported a power relationship between the otolith length and total length of *Micromesistius poutassou* and *Merluccius merluccius* in the Atlantic Ocean while they found a linear relationship in *Trisopterus luscus*, *Trachurus trachurus*, *T. picturatus* and *T. mediterraneus* species. Can (2000) reported that there was a power relationship between the total length-otolith length and fish length-otolith length of *Pagellus erythrinus* in İskenderun Gulf, while this relationship was exponential between total length-otolith weight and linear between otolith length-otolith weight. Bostancı et al. (2009) found a linear relationship between the otolith length and fork length of *Sander lucioperca* samples in Lake Eğirdir. Yılmaz et al. (2014) indicated a non-linear function (power model) between total length-otolith length, total length-otolith height and total length-otolith weight in females and males of *Perca fluviatilis* in Lake Ladik. Yılmaz et al. (2015) investigated the relationships between dimensions (length and width) of the asteriscus and the lapillus otoliths and total length for five cyprinid species (*Abramis brama*, *Blicca bjoerkna*, *Carassius gibelio*, *Chondrostoma regium* and *Scardinius erythrophthalmus*) inhabiting Lake Ladik. They found nonlinear (power) relationships between total length and otolith (asteriscus and lapillus) size in *A. brama*, *B. bjoerkna*, *C. gibelio* and *S. erythrophthalmus* in this lake. The same researchers suggested linear regression was the most appropriate model for *C. regium*.

Casselmann (1979, 1996) stated that cleithrum length was a significant indicator of fish length in Esocid species and the relationships could be defined correctly through linear regression. The same researcher made an important contribution to the practical usage of such relationships by showing that approximately 10 times of the cleithrum length was equal to fish total length. Engstedt et al. (2010) explained the relationship between total length and otolith length in pikes caught from two areas of the Swedish shores of the Baltic Sea by using power model. Our results were similar to Casselman's (1979, 1996) results, yet different from the findings of Engstedt et al. (2010).

The strong relationships were found between the otolith and cleithrum dimensions and fish total lengths of *Esox lucius* in Lake Ladik. In addition, cleithrum length was found to be more convenient

in the back computation of fish length when compared with the different otolith measurements. The data of this study can be used to back-calculated fish length from recovered otolith and cleithrum found in the stomachs of piscivorous animals such as fishes and water birds. We are of the opinion that the regression equations obtained in this study will be useful for the ichthyologist who investigating the feeding habits of top predator and paleontologist who studying about fish fossils.

#### References

- Apaydin Yagci, M., Yılmaz, S., Yazıcıoğlu, O., Polat, N. (2015): The zooplankton composition of Lake Ladik (Samsun, Turkey). *Turkish Journal of Zoology* 39: 652-659.
- Aydin, R., Caltı, M., Coban, M.Z. (2004): Relationships between fish lengths and otolith length in the population of *Chondrostoma regium* (Heckel, 1843) inhabiting Keban Dam Lake. *Pakistan Journal of Biological Sciences* 7: 1550-1553.
- Bostancı, D., Yılmaz, S., Yılmaz, M., Kandemir, Ş., Polat, N. (2009): Eğirdir Gölü'nden sudak (*Sander lucioperca* L., 1758)'nın otolit boyutları-balık boyu ilişkileri ve bazı populasyon parametreleri. *Firat Univ. Journal of Science* 21(1): 9-17. [in Turkish]
- Campana, S.E. (1999): Chemistry and composition of fish otoliths: pathways, mechanisms and applications. *Marine Ecology Progress Series* 188: 263-297.
- Can, M.F. (2000): İskenderun körfezi'ndeki kırmızı mercan (*Pagellus erythrinus* L., 1758) balığında otolit-balık ilişkileri. *Doğu Anadolu Bölgesi IV. Su Ürünleri Sempozyumu* 28-30 Haziran, 31-38. [in Turkish]
- Casselmann, J.M. (1979): The esocid cleithrum as an indicator calcified structure. pp. 249-272. In: Dube, J., Gravel, Y. (eds), *Proc. 10th warmwater workshop, spec. publ. NE Div. Am. Fish. Soc. Publ. By Que. Min. Loisir, de la Chasse and de la Peche, Dir. De la Recherche Faunique, Montreal, Que, Canada.*
- Casselmann, J.M. (1990): Growth and relative size of calcified structures of fish. *Transactions of the American Fisheries Society* 119: 673-688.
- Casselmann, J.M. (1996): Age, growth and environmental requirements of pike. pp. 69-101. In: Craig, J.F. (ed.), *Pike: Biology and Exploitation*. London: Chapman & Hall.
- Das, M. (1994): Age determination and longevity in fishes. *Gerontology* 40: 70-96.
- Dietrich, J.P., Taraborelli, A.C., Morrison, B.J., Schaner, T. (2006): Allometric relationships between size of calcified structures and round goby total length. *North American Journal of Fisheries Management* 26: 926-931.
- Engstedt, O., Stenroth, P., Larsson, P., Ljunggren, L., Elfman, M. (2010): Assessment of natal origin of pike (*Esox lucius*) in the Baltic Sea using Sr: Ca in otoliths. *Environmental Biology of Fishes* 89: 547-555.
- Faust, M.D., Breeggemann, J.J., Bahr, S., Graeb, B.D.S. (2013): Precision and bias of cleithra and sagittal otoliths used to estimate ages of northern pike. *Journal of Fish and Wildlife Management* 4(2): 332-341.
- Gamboia, D.A. (1991): Otolit size versus weight and body-length relationships for eleven fish species of Baja California, Mexico. *Fishery Bulletin* 89(4): 701-706.
- Granoderio, P.J., Silva, M.A. (2000): The use of otoliths and vertebrae in the identification and size-estimation of fish in predator-prey studies. *Cybius* 24(4): 383-393.

- Harvey, J.T., Loughlin, T.R., Perez, M.A., Oxman, D.S. (2000): Relationship between fish size and otolith length for 63 species of fishes from the Eastern North Pacific Ocean. NOAA Technical Report NMFS 150. Washington, DC, USA: NOAA.
- Jawad, L., Sadighzadeh, Z., Al-Busaidi, H. (2012): The relationship between fish length and otolith dimensions of mugilid fish, *Liza kluzingeri* (Day, 1888) collected from the Persian Gulf near Bandar Abbas. *Annales Series Historia Naturalis* 22: 77-82.
- Laine, A.O., Momot, W.T., Ryan, P.A. (1991): Accuracy of using scales and cleithra for aging northern pike from an oligotrophic Ontario Lake. *North American Journal of Fisheries Management* 11: 220-225.
- Mehanna, S.F., Jawad, L.A., Ahmed, Y.A., Abu-el-Regal, M.A., Dawood, D. (2016): Relationships between fish size and otolith measurements for *Chlorurus sordidus* (Forsskål, 1775) and *Hippocampus harid* (Forsskål, 1775) from the Red Sea coast of Egypt. *Journal of Applied Ichthyology* 32: 356-358.
- Neuman, R.M., Willis, D.W., Sammons, S. (1994): Seasonal growth of northern pike (*Esox lucius*) in a South Dakota Glacial Lake. *Journal of Freshwater Ecology* 9(3): 191-196.
- Rydell, J.J., Jolley, J.C., Phelps, Q.E., Willis, D.W. (2008): Northern pike (*Esox lucius*) population characteristics and relations to recruitment in Hackberry Lake, Nebraska. *Transactions of the Nebraska Academy of Sciences* 31: 43-49.
- Scharf, F.S., Yetter, R.M., Summers, A.P., Juanes, F. (1998): Enhancing diet analyses of piscivorous fishes in the Northwest Atlantic through identification and reconstruction of original prey sizes from ingested remains. *Fishery Bulletin* 96: 575-588.
- Tarkan, A.N., Bilge, G., Gaygusuz, Ö., Tarkan, A.S., Gürsoy, Ç., Acıpnar, H. (2007): On the use of otoliths of a ponto-caspian gobiid *Proterorhinus marmoratus* (Pallas, 1814) from Lake İznik (Turkey) in prey-predator studies. *International Journal of Natural and Engineering Sciences* 1(3): 29-33.
- Traynor, D., Moerke, A., Greil, R. (2010): Identification of Michigan fishes using cleithra. *Great Lakes Fish. Comm. Miscellaneous Publication*, 2010-02.
- Tuset, V.M., Lombarte, A., Assis, A.A. (2008): Otolith atlas for the western Mediterranean, north and central eastern Atlantic. *Scientia Marina* 72(Suppl.1): 7-198.
- Yılmaz, S., Yazıcıoğlu, O., Erbaşaran, M., Esen, S., Zengin, M., Polat, N. (2012): Length-weight relationship and relative condition factor of white bream, *Blicca bjoerkna* (L., 1758), from Lake Ladik, Turkey. *Journal of The Black Sea/Mediterranean Environment* 18(3): 380-387.
- Yılmaz, S., Yazıcıoğlu, O., Saygin (Ayaydin), S., Polat, N. (2014): Relationships of otolith dimensions with body length of european perch, *Perca fluviatilis* L., 1758 from Lake Ladik, Turkey. *Pakistan Journal of Zoology* 46(5): 1231-1238.
- Yılmaz, S., Yazıcıoğlu, O., Yazici, R., Polat, N. (2015): Relationships between fish length and otolith size for five cyprinid species from Lake Ladik, Samsun, Turkey. *Turkish Journal of Zoology* 39: 438-446.