

Knowledge gaps and bibliographical revision about descriptions of free-swimming anuran larvae from Brazil

Diogo Borges PROVETE¹, Michel Varajão GAREY^{2,3},
Fernando Rodrigues DA SILVA⁴ and Mainara Xavier JORDANI^{2,3}

1. Programa de Pós-Graduação em Ecologia e Evolução, Laboratório de Ecologia de Insetos, Departamento de Ecologia, Instituto de Ciências Biológicas, Universidade Federal de Goiás – UFG, CP 131, 74001-970, Goiânia, GO, Brasil e-mail: dbprovete@gmail.com

2. Laboratório de Ecologia Animal, Departamento de Zoologia e Botânica, Instituto de Biociências, Letras e Ciências Exatas, Universidade Estadual Paulista-UNESP, R. Cristóvão Colombo, 2265, Jd. Nazareth, 15054-000, São José do Rio Preto, São Paulo, Brasil, www.ibilce.unesp.br

3. Programa de Pós-Graduação em Biologia Animal, Departamento de Zoologia e Botânica, Instituto de Biociências, Letras e Ciências exatas, Universidade Estadual Paulista-UNESP, R. Cristóvão Colombo, 2265, Jardim Nazareth, 15054-000, São José do Rio Preto, São Paulo, Brasil. e-mail: michelgarey@gmail.com, mainaraxj@yahoo.com.br

4. Universidade Federal de São Carlos - UFSCAR, Campus Sorocaba, Rod. João Leme dos Santos Km 110, CEP 18052-780, Sorocaba, SP, Brasil. e-mail: bigosbio@yahoo.com.br
* Corresponding author, D.B. Provete, e-mail: dbprovete@gmail.com

Received: 31. October 2011 / Accepted: 12. May 2012 / Available online: 02. June 2012 / Printed: December 2012

Abstract. Research on tadpole morphology has grown rapidly in recent decades, but many larval anurans remain unknown. We made a literature survey of all studies that describe free-swimming larvae of anuran species occurring in Brazil in order to identify major gaps and provide a searchable bibliographical source about these descriptions. We found that 61.25% of tadpoles of species with free-swimming larvae have already been described. We hope that this paper will foster future tadpole descriptions by providing easy access to the literature.

Key words: literature survey, tadpoles, Anura, herpetology.

Introduction

Traditionally, most studies on anurans have focused on variation and characteristics of the post-metamorphic stage with a few studies dealing with tadpole morphology and ecology (Larson & De Sá 1998). On the other hand, research on tadpoles is increasing rapidly in the last decades (see McDiarmid & Altig 1999) and many unusual larval features have been reported (e.g., Natale et al. 2011). Even so, tadpoles of most tropical anuran species remain unknown. This fact is even more worrisome because amphibians have been declining or even disappearing from the planet at an alarming rate (Stuart et al. 2004). Information on every anuran life stage, from eggs to adults, is paramount to derive proper conservation and management plans. Nonetheless, difficulties in tadpole identification are a major obstacle in anuran surveys and in the development of conservation and management programs. Trying to balance studies between adults and tadpoles is extremely important because: i) the use of larval characters in taxonomic studies can assist in solving phylogenetic issues among species and genera (Larson and De Sá 1998; Frost et al. 2006) and even at the ordinal level (Haas 2003); ii) tadpoles are appropriate

models in a broad range of ecological approaches, such as ecotoxicology (Relyea 2005), competitive and predatory interactions (Morin 1983, 1986), food webs (Wilbur 1997), and community structure (Wilbur 1987); and iii) most tadpoles remain at the breeding sites longer than the adults (Altig & McDiarmid 1999). Therefore, these factors make tadpoles a critical component in any anuran survey (Shaffer et al. 1994, Skelly & Richardson 2010, Silva 2010).

The current estimate of amphibian species in the world is about 6,921 (AmphibiaWeb 2012), but this number is constantly changing, as new species are being discovered either/or becoming extinct. Brazil harbors the greatest species richness of amphibians in the world, with about 877 species (SBH 2010). Among these, 849 are anurans, of which 738 have free-swimming larvae that inhabit a wide variety of habitats (e.g., bromeliads, streams, and ponds; Haddad and Prado 2005).

During a workshop held in 2004, researchers estimated that about 40% of the tadpoles had not been described yet (Andrade et al. 2007). Our goal here is to review the literature about tadpoles of Brazilian species in order to identify major knowledge gaps. In addition, we provide an on-line and fully searchable database that might be helpful for

South American herpetologists, who will have an easy access to the literature about Brazilian free-swimming larvae, including internal oral features and chondrocranial morphology.

Materials and methods

We conducted a literature survey on Internet databases, such as Google scholar, Scielo, Web of Science, and major herpetological journals dealing with descriptions of tadpoles from Brazil. Taxa with direct development (*sensu* Haddad and Prado 2005) were obviously not included (e.g., Brachycephalidae, Craugastoridae, Eleutherodactylidae, some Hemiphractidae, Strabomantidae, and some Microhylidae). Our survey is also based on interviews with other expert herpetologists who have been working with Brazilian tadpoles (see Acknowledgements section). We also included larvae of species that were described with little information a long time ago (e.g., *Ceratophrys aurita*, *Dendropsophus berthalutzae*) and need further detailed descriptions. We did not evaluate the quality of the works or provide further comments. This literature survey did not include exotic species (e.g., *Lithobates catesbeianus*). We followed the taxonomy as given in Frost (2011).

Results and discussion

We found that approximately 61.25% of tadpoles of Brazilian anurans with aquatic free-swimming larvae have been described so far. The family Altophrynidae is composed by a single species (Frost 2011), whose tadpole has not been described. The

family Bufonidae has the least percentage of known tadpoles, since only 31 out of 67 (46.3%) species have been described (Fig. 1). The only family in which 100% of tadpoles were described is Ranidae, but this family has only one species in Brazil. The families with larger number of tadpoles described in Brazil are: Ceratophryidae (80% of 5 species), Leptodactylidae (70.7% of 75), Pipidae (66.7% of 3), Hylidae (64.5% of 341), Leiuperidae (69.6% of 56), Aromobatidae (60% of 20), Microhylidae (52.8% of 36), Hylodidae (54.8% of 42), Centrolenidae (50% of 8), Dendrobatidae (50% of 18), Cycloramphidae (53.8% of 65), and Bufonidae (46.3% of 67 - Fig. 2).

Most Brazilian tadpoles have been described after 1960 (Fig. 2). The 1920s pioneer studies dealing with tadpole morphology published in Brazil (Lutz 1926, 1930, Miranda-Ribeiro 1920, 1923, 1926) only included a picture or a line drawing of larvae in most cases, along with data about size and place where they were found, but no further morphological characteristics. Conversely, recent descriptions often provide more detailed descriptions about external morphology, including internal oral features, and the chondrocranium, which makes it easier to compare and identify tadpoles found in different regions of the country. However, geographical variation in tadpole morphology is scarcely studied in Brazil. An atypical example is the recently described tadpole of *Elachistocleis cesarii* from Rio Grande do Norte, North-




Figure 1. Distribution of the richness of Brazilian anurans with free-swimming larvae (black bars) and the number of tadpoles described per family (gray bars).




Figure 2. Number of tadpole descriptions within recent decades.

eastern Brazil (Magalhães et al. 2012), which had been previously described as *Elachistocleis* sp. in Rossa-Feres and Nomura (2006) from Northwestern São Paulo, Southeastern Brazil (see Provete et al. 2011 for species identification). Apparently there are striking external morphological differences between the two populations (see Magalhães et al. 2012).

We recommend that along with the external morphological descriptions, authors should also describe eggs in detail and the morphology and coloration of hatchlings. Finally, despite the increase in the number of tadpole descriptions in the recent decades, the number of tadpoles described in Brazil is smaller than in Europe and North America, for example.

As a result of this study, we assembled the list of bibliographical references about Brazilian tadpoles, which is available on-line at <http://goo.gl/opquI>. The user can now search the database directly and copy the reference. The database is maintained and continuously updated by DBP.

Acknowledgments. We are grateful to Albertina Lima, Camila Both, Denise de C. Rossa Feres, Janalee Caldwell, Joice Ruggieri, Luís C. Schiesari, Luiz N. Weber, Marcelo Kokubum, Marcelo Menin, Natacha Y. N. Dias, and Pedro Ivo for providing us with literature about the description of some tadpoles. Délio Baêta, Ronald Altig, George Zug, Leandro Alcalde, and Victor Dill read the first versions of this manuscript. Additionally, the authors are grateful for the reviews by T. Vascelos and M. Solé that greatly improved the overall quality of the manuscript. DBP is supported by a doctoral fellowship from CAPES-DS. MVG (2008/50575-1) and MXJ (#2011/05603-0) are supported by fellowships from FAPESP (Cnpq #563075/2010-4).

References

- Alcalde, L., Natale, G.S., Cajade, R. (2006): The tadpole of *Physalaemus fernandezae* (Anura: Leptodactylidae). *Herpetological Journal* 16: 203-211.
- Altig, R., McDiarmid, R.W. (1999): Diversity: familial and generic characterizations. In: *Tadpoles: the biology of anuran larvae*. Edited by R.W. McDiarmid, R. Altig. University of Chicago Press, Chicago. pp. 295-337
- Amphibiaweb: information on amphibian biology and conservation. (2012): Berkeley, California: amphibiaweb. Available: <http://amphibiaweb.org/>. (accessed: feb 15, 2012).
- Andrade, G.V., Eterovick, P.C., Rossa-Feres, D.C., Schiesari, L. (2007): Estudos sobre girinos no Brasil: histórico, conhecimento atual e perspectivas. pp. 127-145. In: Nascimento, L.B., Oliveira, M.E. (eds), *Herpetologia no Brasil II*. Sociedade Brasileira de Herpetologia, Belo Horizonte.
- Frost, D.R., Grant, T., Faivovich, J., Bain, R.H., Haas, A., Haddad, C.F.B., De Sá, R.O., Channing, A., Wilkinson, M., Donnellan, S.C., Raxworthy, C.J., Campbell, J.A., Blotto, B.L., Moler, P., Drewes, R.C., Nussbaum, R.A., Lynch, J.D., Green, D.M., Wheeler, W.C. (2006): The amphibian tree of life. *Bulletin of the American Museum of Natural History* 297: 1-370.
- Frost, D.R. (2011): Amphibian species of the world: an online reference. Version 5.5 (31 January, 2011). Electronic database accessible at <http://research.amnh.org/vz/herpetology/amphibia/> - American Museum of Natural History, New York, USA.
- Haas, A. (2003): Phylogeny of frogs as inferred from primarily larval characters (Amphibia: Anura). *Cladistics* 19:23-89.
- Haddad, C.F.B., Prado, C.P.A. (2005): Reproductive modes in frogs and their unexpected diversity in the Atlantic Forest of Brazil. *Bioscience* 55: 207-217.
- Larson, P.M., De Sá, R.O. (1998): Chondrocranial morphology of *Leptodactylus* larvae (Leptodactylidae: Leptodactylinae): its utility in phylogenetic reconstruction. *Journal of Morphology* 238:287-305.
- Lutz, A. (1926): Observações sobre batrachios brasileiros. Parte I: o gênero *Leptodactylus* Fitzinger. *Memórias do Instituto Oswaldo Cruz* 19: 139-174.
- Lutz, A. (1930). Observações sobre batrachios brasileiros. *Taxonomia e biologia das elossinas*. *Memórias do Instituto Oswaldo Cruz* 24: 195-222.
- Magalhães, F.M., Santana, D.J., Neto, A.M., Garda, A.A. (2012): The tadpole of *Elachistocleis cesarii* Miranda-Ribeiro, 1920 (Anura, Microhylidae). *Zootaxa* 3187: 54-56.
- McDiarmid, R.W., Altig, R. (1999): *Tadpoles: the biology of anuran larvae*. University of Chicago Press, Chicago.
- Miranda-Ribeiro, A. (1920): Os hylodídeos do museu paulista. *Revista do Museu Paulista* 13: 813-846.

- Miranda-Ribeiro, A. (1923): Observações sobre algumas fases evolutivas de *Ceratophrys* e *stombus*. Arquivos do Museu Nacional 24: 201-205.
- Miranda-Ribeiro, A. (1926). Notas para servirem ao estudo dos Gymnobatrachios (Anura) brasileiros. Arquivos do Museu Nacional 27: 7-227.
- Morin, P.J. (1983): Predation, competition, and the composition of larval anuran guilds. Ecological Monographs 53:119-138.
- Morin, P.J. (1986): Interactions between interspecific competition and predation in an amphibian predator-prey system. Ecology 67: 713-720.
- Natale, G.S., Alcalde, L., Herrera, R., Cajade, R., Schaefer, E.F., Marangoni, F., Trudeau, V.L. (2011): Underwater acoustic communication in the macrophagic carnivorous larvae of *Ceratophrys ornata* (Anura: Ceratophryidae). Acta Zoologica 92: 46-53.
- Provete, D.B., Garey, M.V., Silva, F.R., Rossa-Feres, D.C. (2011): Anurofauna do noroeste paulista: lista de espécies e chave de identificação para adultos. Biota Neotropica 11: 377-391.
- Relyea, R.A. (2005): The impact of insecticides and herbicides on the biodiversity and productivity of aquatic communities. Ecological Applications 15: 618-627.
- Rossa-Feres, D.C., Nomura, F. (2006): Characterization and taxonomic key for tadpoles (Amphibia: Anura) from the northwestern region of São Paulo State, Brazil. Biota Neotropica 6(1): dx.doi.org/10.1590/S1676-06032006000100014.
- SBH. (2010). Brazilian amphibians - list of species. <http://www.sberpetologia.org.br>. <Last accessed 7 July 2010>.
- Shaffer, H.B., Alford, R.A., Woodward, B.D., Richards, S.J., Altig, R., Gascon, C. (1994): Quantitative sampling of amphibian larvae. pp. 130-140. In: Heyer, W.R. et al. (eds), Measuring and monitoring biological diversity - standard methods for amphibians. Washington, Smithsonian Institution Press.
- Silva, F.R. (2010): Evaluation of survey methods for sampling anuran species richness in the neotropics. South American Journal of Herpetology 5: 212-220.
- Skelly, D.K., Richardson, J.L. (2010): Larval sampling. pp. 55-70. In: Dodd Jr., C.K. (ed.), Amphibian ecology and conservation - a handbook of techniques. Oxford, Oxford University Press.
- Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L., Waller, R.W. (2004): Status and trends of amphibian declines and extinctions worldwide. Science 306: 1783-1786.
- Wilbur, H.M. (1987): Regulation of structure in complex system: experimental temporary pond communities. Ecology 68: 1437-1452.
- Wilbur, H.M. (1997): Experimental ecology of food webs: complex system in temporary ponds. Ecology 78: 2279-2302.
-